

## NPY1 Receptors Participate in the Regulation of Myocardial Contractility in Rats

Masliuko P., Anikina T., Zverev A., Krylova A., Moiseev K., Zefirov T.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

### Abstract

© 2017, Springer Science+Business Media New York. Selective agonist (Leu(31)Pro(34)NPY) and blocker (BIBP-3226) of NPY1 receptors were used to determine the type of NPY receptors involved in myocardial contraction. Experiments with isometric contraction of myocardial strips from mature rats showed that the agonist produced the most potent effect in a concentration of  $10^{-7}$  M. In this concentration, Leu(31)Pro(34)NPY showed the greatest positive inotropic effect on the contraction of the atria and ventricles. In contrast, selective blocker BIBP-3226 reduced the force of myocardial contractions. Pretreatment of myocardial strips with this blocker abolished the positive inotropic effect of Leu(31)Pro(34)NPY, which attested to important role of NPY1 receptors in myocardial contraction.

<http://dx.doi.org/10.1007/s10517-017-3629-x>

---

### Keywords

myocardial contractility, neuropeptide Y, rat, types of NPY receptors

### References

- [1] Zverev AA, Anikina TA, Maslyukov PM, Zefirov TL. Role of neuropeptide Y in myocardial contractility of rats during early postnatal ontogeny. *Bull. Exp. Biol. Med.* 2014;157(4):421-423.
- [2] Nozdachev AD, Masliukov PM. Neuropeptide Y and autonomic nervous system. *J. Evol. Biochem. Physiol.* 2011; 47(2):121-130.
- [3] Bischoff A, Michel MC. Renal effects of neuropeptide Y. *Pflugers Arch.* 1998;435(4):443-453.
- [4] Masliukov PM, Moiseev K, Emanuilov AI, Anikina TA, Zverev AA, Nozdachev AD. Development of neuropeptide Y-mediated heart innervation in rats. *Neuropeptides.* 2016;55:47-54.
- [5] Mollereau C, Gouardères C, Dumont Y, Kotani M, Detheux M, Doods H, Parmentier M, Quirion R, Zajac JM. Agonist and antagonist activities on human NPFF(2) receptors of the NPY ligands GR231118 and BIBP-3226. *Br. J. Pharmacol.* 2001;133(1):1-4.
- [6] Protas L, Robinson RB. Neuropeptide Y contributes to innervation-dependent increase in I(Ca, L) via ventricular Y2 receptors. *Am. J. Physiol.* 1999;277(3, Pt 2):H940-H946.
- [7] Wahlestedt C, Wohlfart B, Hakanson R. Effects of neuropeptide Y (NPY) on isolated guinea-pig heart. *Acta Physiol. Scand.* 1987;129(4):459-463.